

Read the label before opening the container.

For full particulars, see enclosed leaflet.

Aceto 900



South Africa Reg. No: L 7863 Act No. 36 of 1947

Namibia Reg. No: N-AR 1196 Act no 36 of 1947

An emulsifiable concentrate herbicide for the pre-emergence control of grasses and certain broadleaf weeds in ratoon sugarcane and Eucalyptus and pine plantations and post-emergence control in ratoon sugarcane.

HRAC HERBICIDE GROUP CODE: K3

ACTIVE INGREDIENT:

Acetochlor (acetanilide)..... 900g/ℓ

Product Information Tel no: 072 678 8226

In case of poisoning: 082 446 8946

www.envirobiochem.co.za

Expiry Date:

Batch No:

Date of Manufacture:

UN No. 3082

enviro
bio-chem

Registration holder:

Erintrade CC t/a RT Chemicals

Reg. No: CK 2001/036403/23

44 Kerk Street, Lichtenburg

North West, South Africa 2740

Tel: +27 87 231 7261



CAUTION
VERSIGTIG



WARNINGS:

- Poisonous when swallowed.
- May cause eye and/or skin irritation.
- Toxic to fish.
- Keep out of reach of children, uninformed persons and animals.
- Store in the original container under lock and key, away from food, feedstuffs, fertilizers and seed.
- Store in a cool dry place.
- **Re-entry:** Do not enter the treated field until the spray deposit has dried unless wearing protective clothing.

Although this herbicide has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be effected by factors such as abnormal soil, climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well as by the method, time and accuracy of the application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the herbicide concerned due to failure of the user to follow the label instructions or to the occurrence of conditions that could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS:

- Do not breathe in fumes or spray mist.
- Wear rubber gloves when handling the concentrate.
- After use and in case of accidental skin contact, wash thoroughly with soap and water.
- In the case of accidental eye contact, rinse the eyes with clean water for at least 15 minutes. Get medical attention if necessary.
- Do not eat, drink or smoke whilst mixing or applying or before washing hands and face and change of clothing after use.
- Prevent contamination of food, feedstuff, eating utensils and drinking water.
- Prevent drift of spray mist onto other crops, grazing, rivers, dams or areas not under treatment.
- Rinse empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy empty container by perforation and flattening and never use for any other purpose.
- Clean applicator after use and dispose of wash water where it will not contaminate crops, grazing, dams or rivers.
- Low temperatures and/or heavy prolonged rains soon after planting may cause damage to the crop.

RESISTANCE WARNING:

For resistance management **Aceto 900** is a **group code K3** herbicide. Any weed population may contain individuals naturally resistant to **Aceto 900** and other **group code K3** herbicides. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. **Aceto 900** or any other **group code K3** herbicide may not control these resistant weeds.

To delay herbicide resistance:

- Avoid the exclusive repeated use of herbicides from the same herbicide group code.
- Alternate or tank mix with products from different herbicide group codes.
- Integrate other control methods (chemical, cultural, biological) into weed control programmes.

For specific information on resistance management contact the registration holders of this product.

DIRECTIONS FOR USE: Use only as indicated.

General Information:

- Adhere to restrictions on rotation crops on the labels of herbicides mixed with **Aceto 900**. Before using **Aceto 900** in combination with other herbicides, the labels must be read and recommendations and restrictions adhered to.
- **Aceto 900** must not be applied to poorly drained soils as the herbicide may cause crop injury in combination with waterlogging.
- **Aceto 900** must not be applied to sandy soils that are susceptible to wind erosion.

- Flood irrigation can reduce weed control performance.
- Optimum weed control can be obtained on a fine even seedbed, free of weeds, trash and clods. **Aceto 900** will give weed control in stubble mulch or reduced tillage conditions.

Crop Restrictions – Maize

Aceto 900 and **Extrazine 500 SC** should be applied as a split application only (not in a tankmix). Do not use this combination on the low potential soils of the North West Province, North and Western Free State.

Mixing Instructions:

- Shake the container before use.
- Mixing sequence:
 1. Fill the spray tank to three-quarters full with clean water.
 2. Add the required quantity of complementary herbicide.
 3. Agitate during mixing and spraying.
 4. Add the required **Aceto 900** volume.
 5. Fill the spray tank to the required volume.

Do not allow spray mixes to stand in the spray tank.

Application Information:

- Spray equipment must be accurately calibrated and checked regularly before and during the spraying operation.
- **Aceto 900** alone and in tank mixes must be applied in maize, groundnuts and sugarcane preferably with planting or immediately after planting. Pre-emerge treatment must not be applied on maize later than three days after planting and/or the last cultivation.
- Use 100 to 300ℓ/ha spray volume for overall ground application.
- Spray volumes and dosage rates should be adjusted accordingly for on the row band treatments.
- 10 to 15 mm rain within 7 to 10 days after application for good weed control is necessary.
- For optimum weed control seedbed preparation should take place within 3 days before planting. Planting is followed immediately with the **Aceto 900** application and then 10 to 15mm rain or irrigation within 3 days to wash the herbicide into the zone of germinating weeds.
- If conditions are dry, weed seedlings may emerge. These can be controlled with either a shallow cultivation (2 to 4cm deep) or with a recommended post-emerge herbicide treatment.
- If soil crusting becomes a problem, use a rotary harrow in the same direction as the planted rows to assist maize emergence.
- After application harrowing may reduce weed control if untreated soil is brought to the surface.
- Sufficient fertilizer must be placed near the seed when planting to promote vigorous seedling growth. This is very important with early planting when the soil temperature is relatively low and is accompanied by continuous moist conditions.

WEEDS CONTROLLED BY Aceto 900

Sweet signal grass	<i>Brachiaria eruciformis</i>
Feathertop chloris	<i>Chloris virgata</i>
Crab finger grass	<i>Digitaria sanguinalis</i>
Goose grass	<i>Eleusine indica</i>
Common buffalo grass	<i>Panicum maximum</i>
Sweet buffalo grass	<i>Panicum schinzii</i>
Garden brittle grass	<i>Setaria pallide-fusca</i>
Herringbone grass	<i>Urochloa panicoides</i>
Perennial pigweed	<i>Amaranthus deflexus</i>
Common pigweed	<i>Amaranthus hybridus</i>

WEEDS CONTROLLED BY Aceto 900

Thorny pigweed Red pigweed White goosefoot Green goosefoot Bengal wandering Jew Thorn apple (early germinating only) Bladder weed Wild gooseberry Purslane Dwarf marigold Khaki weed (early germinating only)	<i>Amaranthus spinosus</i> <i>Amaranthus thunbergii</i> <i>Chenopodium album</i> <i>Chenopodium carinatum</i> <i>Commelina benghalensis</i> <i>Datura spp</i> <i>Hibiscus trionum</i> <i>Physalis angulata</i> <i>Portulaca oleracea</i> <i>Schkuhria pinnata</i> <i>Tagetes minuta</i>
---	---

Since Aceto 900 controls annual grasses, other annual grasses that are not listed may be controlled to a greater or less degree. The registration holder does not accept liability in respect of the control of unlisted weeds.

SUGARCANE: PRE-EMERGENCE APPLICATION

Aceto 900 can be used in combination with other herbicides for pre- and post-emerge weed control in ratoon sugarcane.

SUGARCANE TABLE 1

Annual grasses, broadleaf weeds and *Cyperus esculentus* (yellow nutsedge) control.

Aceto 900 2 - 3ℓ/ha

In a tank mix with one of;

Or AMETRYN 500 SC (Reg. No: 8219) 2 - 3ℓ/ha

Or Diuron 800 SC 3 ℓ/ha

Or Atrazine 500 SC 2 - 5ℓ/ha (0 - 35% clay)

Or Atrazine 500 SC 3 - 5ℓ/ha (>35% clay)

The higher dosage rates should be used on heavier soils (>30% clay) and also where a longer residual action is required and/or for improved control of *Cyperus esculentus* (yellow nutsedge). The lower dosage rates should be used on lighter soils.

SUGARCANE: POST-EMERGENCE APPLICATION

SUGARCANE TABLE 2

Annual grasses, broadleaf weeds and *Cyperus esculentus* (yellow nutsedge) control.

REMARKS

APPLICATION ℓ/ha

After the 5-leaf stage of the cane the treatment should be directed between the rows.

Aceto 900 2 - 3

+ AMETRYN 500 SC (Reg. No: 8219) 6

+ BOOSTER OIL (Reg. No: L 6148)

May only be applied up to and including the 2-3 leaf stage of the sugarcane.

Controls *Cyperus esculentus* (yellow nutsedge).

OR

Aceto 900 2 - 3

+ AMETRYN 500 SC 3 - 5

+ PARAQUAT SL (Reg. No: 7650) 1.5

SUGARCANE: POST-EMERGENCE APPLICATION

SUGARCANE TABLE 2

Annual grasses, broadleaf weeds and *Cyperus esculentus* (yellow nutsedge) control.

REMARKS	APPLICATION ℓ/ha
After the 5 leaf stage of the sugarcane the treatment should be directed between the rows. Controls <i>Cyperus esculentus</i> (yellow nutsedge).	Or Aceto 900 2 - 3 + Diuron 800 SC 2.5 - 3 + SPRITE 255 (Bromoxynil 255 EC) (Reg. No: 8247) 1.25
May only be applied up to and including the 3 leaf stage of the sugarcane. Controls <i>Cyperus esculentus</i> (yellow nutsedge).	Or Aceto 900 2 - 3 + Diuron 800 SC 2.5 – 3 + PARAQUAT SL 1.5
May only be applied up to and including the 2 - 3 leaf stage of the sugarcane. Controls <i>Cyperus esculentus</i> (yellow nutsedge).	Or Aceto 900 2 - 3 + Atrazine 500 SC 2 - 5 (0-35% clay) Or + Atrazine 500 SC 3 - 5 (>35% clay)
<ul style="list-style-type: none"> • The higher dosage rates should be used on heavier soils. • All post-emergence treatments should be made before tillering of annual grasses. • Aceto 900 will not cause foliar injury to sugarcane. However, certain herbicides in the above tank mix combinations may cause injury. Consult the relevant labels regarding use restrictions prior to use. 	

PLANTATIONS

Eucalyptus and Pine Plantations

Spray existing weeds with KLEEN UP (Reg. No: 5853). Follow this with Aceto 900 as a pre-plant treatment or within 3 days after transplanting young trees.

Use the lower rate for annual grass control.

Use a minimum of 1.5ℓ/ha for *Digitaria sanguinalis* (crab finger grass) control.

Use the higher rates for extended residual control and/or improved *Cyperus esculentus* (yellow nutsedge) control.

CLAY %	Aceto 900 ℓ/ha
0 - 10	0.75 - 1.5
11 - 30	1 - 3