

Read the label before opening the container.

For full particulars, see enclosed leaflet.

Booster OIL

South Africa Reg. No: L 6148 Act No. 36 of 1947

An emulsifiable vegetable oil for use as a wetter, sticker, spreader.

ACTIVE INGREDIENT:

Vegetable oil851g/l

Product Information Tel no: 072 678 8226

In case of poisoning: 082 446 8946

www.envirobiochem.co.za

Expiry Date:

Batch No:

Date of Manufacture:

The logo for enviro bio-chem features the word "enviro" in a bold, lowercase sans-serif font, with a green leaf icon above the letter "i". Below "enviro" is a horizontal line, and under that line, the words "bio-chem" are written in a smaller, lowercase sans-serif font.

Registration holder:

RT Chemicals CC

Reg. No: CK 87/21925/23

44 Kerk Street, Lichtenburg

North West, South Africa 2740

Tel: +27 87 231 7261



CAUTION
VERSIGTIG



WARNINGS:

- Handle with care.
- Observe all warnings that are applicable to products with which **Booster OIL** will be used.
- Store away from food and feedstuffs.
- Keep out of reach of children, uninformed persons and animals.

Although this remedy has been extensively tested under a wide variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

- Observe all precautions that are applicable to products with which **Booster OIL** will be used.
- Do not inhale the spray mist.
- After use and in case of accidental skin contact, wash thoroughly with soap and water.
- In case of eye contact flush eyes with copious amounts of clean water for at least 15 minutes.
- Wash contaminated clothing daily.
- Do not eat, drink or smoke while mixing or applying or before washing hands and face and change of clothing.
- Prevent drift of spray mist onto other crops, grazing, rivers, dams or areas not under treatment.
- Thoroughly clean applicator after use and dispose of wash water where it will not contaminate crops, grazing, rivers or dams.
- Triple rinse empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy empty container by perforation and flattening and never use for any other purpose.
- Prevent contamination of food, feedstuffs, drinking water and eating utensils.

DIRECTIONS FOR USE: Use only as indicated.

Compatibility:

Booster OIL is compatible with most pesticides but it is advisable to do small scale compatibility tests prior to mixing large quantities.

Mixing Instructions:

- **Booster OIL** must always be premixed with pesticides/and or foliar sprays prior to adding to the spray tank and water [except soluble concentrates – see below].
- For wettable powder or dispersible granular formulations, add water to the pesticide in equal amounts and mix into a slurry. Add this to the required amount of **Booster OIL** (premix) and follow the mixing instructions.
- Fill the spray tank with clean water to half the required volume and start agitation. Add the premix to the spray tank. Top up the spray tank to the required spray volume with water while continuously agitating.
- For soluble concentrates, fill the spray tank with clean water to half the required volume and add the required volume of the pesticide. Agitate the contents of the spray tank and add the required volume of **Booster OIL**. Top up the spray tank to the required volume with water while continuously agitating.

APPLICATION RATES:

APPLICATION METHOD	RATES
Aerial application - aeroplane:	250m l - 2l/ha
Mistblower application	1 - 2l/ha (low volume)
Handgun application	200mℓ/100ℓ water (spot spray)
Knapsack sprayer application	20 - 30mℓ/10ℓ water
Tree orchard application	150 - 200mℓ/100ℓ water of 1.5 - 2.5ℓ/ha

- NOTE:**
1. Use the higher rate when using large dosage rate products.
 2. Avoid spray drift and particularly spraying when conditions are conducive to spray drift.