

Read the label before opening the container.

For Full particulars, see enclosed leaflet.

EmGee



South Africa Reg.No: L 8029 Act No. 36 of 1947

A wettable granule post-emergence herbicide packed in water soluble sachets
for the control of broadleaf weeds in soy beans as indicated.

HRAC HERBICIDE GROUP CODE: B

ACTIVE INGREDIENT:

Chlorimuron-ethyl (sulfonyl urea)250 g/kg

Product Information Tel no: 082 801 6759
In case of poisoning: 082 446 8946

www.enviro-crop.co.za

enviro
efficiency • quality • competitiveness

Batch No:
Date of Manufacture:

Registration holder:

Erintrade cc t/a RT Chemicals
CK 2001/036403/23
10 Ninth Avenue, Industria, Kroonstad, 9500
Tel: 056 213 1967

06082012-10/12



CAUTION
VERSIGTIG

WARNINGS

- Allow a minimum of 60 days between application and harvest of soy beans.
- Direct exposure to the product may irritate eyes, skin and mucous membranes and may result in general malaise.
- Keep out of reach of children, uninformed persons and animals.
- Store in the original container in a cool dry place.
- Store away from food and feeds
- Damage to the crop may occur if the recommended dosage is exceeded.
- During periods of lengthy wet, cold conditions soon after **EmGee** application, a yellowing of the crop leaves and, in extreme cases, temporary stunting may occur.
- **Re-entry:** Do not enter the treated field until the spray deposit has dried unless wearing protective clothing.
- **Aerial Application:** Notify all inhabitants in the immediate vicinity of the area to be sprayed and issue the necessary warnings. Do not spray over or allow drift to contaminate water or adjacent areas. Should the recommended dosage rate be exceeded, damage to crops may occur.

Although this herbicide has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be effected by factors such as abnormal soil, climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the weed against the herbicide concerned as well as by the method, time and accuracy of the application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the herbicide concerned due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS

- Do not inhale the dust, fumes or spray mist of this product.
- Wear protective clothing when handling the concentrate (eye protection, rubber boots and gloves).
- In case of accidental contact with eyes, rinse eyes with clean water for at least 15 minutes.
- After use and in case of accidental skin contact, wash the skin thoroughly with soap and water.
- Wash contaminated clothing after use.
- Do not eat, drink or smoke whilst mixing or applying or before washing hands and face and change of clothing after use.
- Prevent contamination of food, feedstuff, eating utensils and drinking water.
- Prevent drift of spray mist onto other crops, grazing, rivers, dams or areas not under treatment.
- Clean the applicator after use and dispose of wash water where it will not contaminate crops, grazing, dams or rivers. After use, rinse the spray equipment with a 1% solution of household ammonia. Allow this solution to stand in the spray equipment for at least 10 minutes (tank, pipes, nozzles). Repeat the rinsing with the ammonia solution before rinsing the equipment twice with clean water.
- Rinse the empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the contents of the spray tank before destroying the container in the prescribed manner. Destroy empty container by perforation and flattening and never use for any other purpose.

RESISTANCE WARNING

For resistance management **EmGee** is a group code B herbicide. Any weed population may contain individuals naturally resistant to **EmGee** and other group code B herbicides. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. **EmGee** or any other group code B herbicide may not control these resistant weeds. To delay herbicide resistance:

- Avoid exclusive repeated use of herbicides from the same herbicide group code.
- Alternate or tank mix with products from different herbicide group codes. Integrate other control methods (chemical, cultural, biological) into weed control programmes.
- For specific information on resistance management contact the registration holder of this product.

USE RESTRICTIONS:

- **Follow-up crops:**
 - Soy beans may be planted at any time after an **EmGee** application.
 - Do not plant wheat within 3 months of an **EmGee** application.
 - Do not plant any other crop within 9 months of an **EmGee** application.
 - The extended residual activity of **EmGee** to sensitive follow-up crops may result from (i) application to soils with pH exceeding 7 (water) and/or (ii) soils containing free lime or (iii) where more than one **EmGee** application is made per season. Under these circumstances consult your technical advisor for specific recommendations.
- Soil factors such as pH, moisture and temperature may influence weed growth and affect the efficacy of **EmGee**.
- The efficacy of **EmGee** will be adversely affected if cool, dry conditions prevail after spraying.
- The spray mixture must be used immediately after mixing. Agitate the mixture continuously and do not allow it to stand for any period of time.
- Do not apply **EmGee** if rain is expected within 1 hour.
- Do not apply **EmGee** before the third soy bean trifoliolate (leaf) opens or later than 60 days before soybean maturity.
- Temporary leaf yellowing and/or leaf crinkling may occur to soy beans 5 to 7 days after application of **EmGee**. These symptoms are associated together with other stress factors. The crop will recover under favourable growing conditions and yield will not be affected.
- Do not apply **EmGee** to plants under stress. Do not apply to plants suffering from heat or cold stress, drought or water-logged soil, disease or insect or other herbicide injury, as crop injury or poor weed control may result. Severe stress such as drought, disease, insect damage or nutrient deficiency following application may also result in crop injury. Delay application until stress conditions are no longer a detrimental factor and weeds and soy beans resume normal growth.
- Do not tank mix **EmGee** with organophosphate insecticides.
- Do not apply **EmGee** more than twice in a growing season to the same soy bean crop.

DIRECTIONS FOR USE Use only as directed.

COMPATIBILITY

Do not mix **EmGee** with products other than those recommended on this label.

MIXING INSTRUCTIONS

Mix the required amount of **EmGee** with a small amount of water to form a smooth paste.

Half fill the spray tank with clean water.

Add the pre-mixed paste to the tank through a sieve and fill the tank with water to the required volume whilst agitating continuously.

Add the required quantity of non-ionic surfactant whilst agitating continually and apply when thoroughly mixed.

Note: Ensure that thorough agitation of the spray mixture is maintained at all times.

APPLICATION

Ground Application:

Apply with a tractor mounted boom sprayer calibrated to spray 200 to 300 ℓ spray mixture per hectare and to give an even spray deposit on the target weeds. Avoid overlapping of spray swaths and prevent over-application.

Aerial Application:

Aerial application may only be done by a registered aerial application operator using a registered and correctly calibrated aircraft according to the instructions of SABS Code 10118 (Aerial Application of Agricultural Remedies). It is important to ensure that the spray mixture is distributed evenly over the target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria be met:

Equipment

- Use whirljet nozzles to produce the desired droplet and coverage of the target area and that will ensure the minimum loss of product through drift.
- Use a nozzle set-up that will produce a droplet spectrum with the lowest possible relative span.
- All nozzles and atomisers should be positioned within the inner 75% of the wingspan to prevent droplets from entering the wingtip vortices.

Application Parameters

- A minimum volume of 30 ℓ spray mixture per ha is recommended. As this product has not been evaluated at a reduced volume rate, the registration holder cannot guarantee efficacy nor be held responsible for any adverse effects if this product is applied aerially at a lower volume rate than recommended.
- A droplet coverage of 30 to 40 droplets per square cm must be recovered on the target
- A droplet spectrum with a VMD of 250 to 300 microns is recommended.
- The height of the spray boom should be maintained at between 3 and 4 m above the target.
- Do not spray when the aircraft is in a climb, at the top of the turn or during a dive or when banking.
- Apply before the crop growth becomes too dense, which will interfere with overall plant coverage.

Meteorological Conditions

- The difference between the wet and dry bulb readings as determined by a whirling hygrometer must not exceed 8°C.
- Do not spray under turbulent, unstable conditions or during the heat of the day when rising thermals and downdraughts occur.
- Do not spray under temperature inversion conditions (spraying in or above the inversion layer).
- Do not spray when the wind speed exceeds 15 km per hour.

General

- Ensure that fields are accurately marked and that the aerial spray operator knows exactly which fields to spray.
- Obtain an assurance from the aerial spray operator that the above requirements will be met.
- Do not spray in the heat of the day.

DOSAGE RATE: Soy beans

CROP	DOSAGE	REMARKS
Soy beans	35 g per ha	Apply to young actively growing weeds. Do not apply to weeds suffering from stress. Add a non-ionic surfactant to EmGee before spraying. Use Booster Oil (L 6148, Act No. 36 of 1947) at a concentration of 0,1 to 0,2% (100 to 200 mL per 100 litres spray solution).

NOTE: A second application 14 to 21 days later may be necessary to control newly germinated weeds which were not adequately controlled by the first application.

WEEDS CONTROLLED:

The following weeds are normally controlled by **EmGee** provided that the following spraying stages (leaf growth stages) are complied with:

COMMON NAME	BOTANICAL NAME	MAXIMUM LEAF STAGE AT APPLICATION
Cape pigweed	<i>Amaranthus hybridus</i>	4
Red pigweed	<i>Amaranthus thunbergii</i>	4
Black jack	<i>Bidens pilosa</i>	4
Swine Cress	<i>Coronopus didymus</i>	4
Large thorn apple	<i>Datura ferox</i>	4
Thorn Apple	<i>Datura stramonium</i>	4
Tumbleweed	<i>Leucas martinicensis</i>	4
Apple of Peru	<i>Nicandra physaloides</i>	4
Wild radish	<i>Raphanus raphanistrum</i>	4
Mexican clover	<i>Richardia brasiliensis</i>	4
Black nightshade	<i>Solanum nigrum</i>	4
Tall khaki weed	<i>Tagetes minuta</i>	4
Klitsbossie	<i>Triumfetta species</i>	4
Cocklebur	<i>Xanthium strumarium</i>	6

VARIABLE WEED CONTROL

COMMON NAME	BOTANICAL NAME
Wandering Jew	<i>Commelina benghalensis</i>
Yellow nutsedge	<i>Cyperus esculentus</i>
Morning Glory	<i>Ipomoea purpurea</i>
Common Dubbeltjie	<i>Tribulis terrestris</i>