

Read the label before opening the container.

For full particulars, see enclosed leaflet.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

BROMACIL 800 WP

South Africa Reg. No: L7934 Act No. 36 of 1947

HRAC HERBICIDE GROUP CODE: C

A wettable powder herbicide with long residual weed control for use in industrial areas.

ACTIVE INGREDIENT:

Bromacil (uracil).....800g/kg

Product Information: 072 678 8226
In case of poisoning: 082 446 8946

HAZARD STATEMENTS

- Harmful if swallowed
- Very toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENTS

- Wash hands, forearms and face thoroughly after handling.
- Dispose of contents in accordance with local/regional/national regulations.



enviro
bio-chem

Registration holder: **Erintrade CC t/a RT Chemicals**

Reg No: CK2001/036403/23

44 Kerk Street, Lichtenburg North West, South Africa 2740

Tel: +27 18 632 0050 | Fax: +27 86 541 7948

Web: www.envirobiochem.co.za

Expiry Date:

Batch No:

Date of Manufacture:

UN No. 3077

WARNINGS:

- Harmful if swallowed
- Very toxic to aquatic life with long lasting effects.
- Toxic to fish, bees and wildlife.
- Store in a cool place away from food, feeds, seed, fertilizers and other agricultural chemicals.
- Keep out of reach of children, uninformed persons and animals.
- Re-entry period: Do not enter the treated area until the spray deposit has dried unless wearing protective clothing.

Although this herbicide has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be effected by factors such as abnormal soil, climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the weed against the herbicide as well as by the method, time and accuracy of the application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS:

- Wash hands, forearms, and face thoroughly after handling
- Do not eat, drink, or smoke when using this product.
- Prevent contamination of food, feedstuffs, drinking water and eating utensils and wash contaminated clothing daily.
- Prevent drift of spray mist onto crops, grazing, rivers, dams or areas not under treatment.
- Clean applicator after use and dispose of wash water where it will not contaminate food, grazing, rivers or dams.
- Triple rinse empty container in the following manner: Invert the empty container over the spray or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the content of the spray tank. Empty containers should be taken for local recycling or waste disposal. Do not re-use empty containers.
- Collect spillage.

RELEVANT SUBSTANCES:

Bromacil 84%

Sodium lauryl sulphate <5%

FIRST AID TREATMENT:

INGESTION: Immediately call a POISON CENTER or doctor/physician.

SKIN CONTACT: (or hair) Remove/Take off immediately all contaminated clothing. Rinse skin with water/shower.

INHALATION: Remove person to fresh air and keep comfortable for breathing.

EYE CONTACT: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do – continue rinsing.

RESISTANCE WARNINGS:

For resistance management **BROMACIL 800 WP** is a **group code C1** herbicide. Any weed population may contain individuals naturally resistant to **BROMACIL 800 WP** and other **group code C1** herbicides. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. **BROMACIL 800 WP** or any other **group code C1** herbicide may not control these resistant weeds.

To delay herbicide resistance:

- Avoid exclusive repeated use of herbicides from the same herbicide group code.
- Alternate or tank mix with products from different herbicide group codes.
- Integrate other control methods (chemical, cultural, biological) into weed control programmes.

For specific information on resistance management contact the registration holders of this product.

USE RESTRICTIONS:

Do not apply near desirable plants or trees such as wind breaks in orchards and ornamental trees along roads and streets or on areas, where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.

Do not use on soil where susceptible crops or any susceptible plants are to be grown in the foreseeable future.

DIRECTIONS FOR USE: Use only as directed.

- Use a sprayer, which will distribute the required amount evenly over the area to be treated.
- Use a coarse spray to reduce drift.
- Best control will be obtained when weeds are short or when excess growth is mowed and removed before application.
- Best results will be obtained if applied shortly before or at the time when the weeds are growing actively.
- Rain or thorough watering after application will cause the material to move to the root zone where it is taken up and thus speed up control.
- The effectiveness and length of time that control will last will depend on the type of soil, amount of rain and other conditions.
- Adequate agitation is essential during mixing and spraying.

SOIL TYPE	DOSAGE/ha	REMARKS
INDUSTRIAL AREAS: Railways, road shoulders, etc.		
<20% Clay	8 tot 10kg	Use the higher rate where existing weeds are treated. Where dense growth of weeds occur a suitable contact herbicide may be included in the spray.
20 tot 35% Clay	10 tot 12kg	
>35% Clay	12 tot 15kg	
Ash	15kg	

Some weed species normally controlled by BROMACIL 800 WP

Upright Starbur	<i>Acanthospermum Hispidum</i>
Cape Pigweed	<i>Amaranthus Hybridus</i>
Tassel Three-Awn	<i>Aristida Congesta</i>
Sweet Signal Grass	<i>Brachiaria Eruciformis</i>
Spiderweb Chloris	<i>Chloris Radiata</i>
Tall Fleabane	<i>Conyza Albida</i>
Common Quick Grass	<i>Cynodon Dactylon</i>
Yellow Nutsedge	<i>Cyperus Esculentus</i>
Purple Nutsedge	<i>Cyperus Rotundus</i>
Crabfinger Grass	<i>Digitaria Sanguinalis</i>
Goose Grass	<i>Eleusine Indica</i>
Common Thatching Grass	<i>Hyparrhenia Hirta</i>
Spotted Cat's Ear	<i>Hypochoeris Radicata</i>
Sweet Buffalo Grass	<i>Panicum Schinzii</i>
Kikuyu Grass	<i>Pennisetum Clandestinum</i>
Purslane	<i>Portulaca Oleracea</i>
Bur Bristle Grass	<i>Setaria Verticillata</i>
Tall Khaki Grass	<i>Tagetes Minuta</i>
Large Carrotseed Grass	<i>Tragus Racemosus</i>