

Read the label before opening the container.

For full particulars, see enclosed leaflet.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

THIS REMEDY CONTAINS AN ANTI-COAGULENT THAT MAY CAUSE INTERNAL BLEEDING IF CONSUMED. THIS REMEDY MAY ONLY BE USED IN TAMPER-RESISTANT BAIT STATIONS AND ONLY AS DIRECTED ON THIS LABEL. OBTAIN, READ AND FOLLOW ALL SAFETY INSTRUCTIONS BEFORE USE.

TORNADO

South Africa Reg. No: L4253 Act No. 36 of 1947

WAX BLOCK BAIT

A wax block bait for the control of roof and Norway rats and house mice, in and around industrial premises as well as rats and mice in newly established plantations

ACTIVE INGREDIENT:

Difenacoum (anti-coagulant).....0.1g/kg

Product Information: 072 678 8226

In case of poisoning: 082 446 8946

Not classified according to GHS

PRECAUTIONARY STATEMENTS

- Obtain, read and follow all safety instructions before use.
- Wear protective gloves.
- If exposed or concerned, get medical advice.
- Store locked up.
- Dispose of contents/container in a responsible manner to avoid accidental exposure of uninformed persons and animals.

enviro
bio-chem

Registration holder: RT Chemicals C.C.

Co. Reg. No: CK87/21925/23

44 Kerk Street, Lichtenburg NW, South-Africa

Tel: +27 18 285 1291 | Fax: +27 86 541 7948

Web: www.envirobiochem.co.za

Expiry Date:

Batch No:

Date of Manufacture:

UN No. 3027

WARNINGS:

- Handle with care.
- Toxic to fish, wildlife and pets.
- Store under lock and key in a cool place away from food and feeds.
- Keep out of reach of children, uninformed persons, pets, poultry and livestock.
- **FLAMMABLE** – Do not store or handle near open flame.
- In case of poisoning call a doctor and make this label available to him.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS:

- Obtain, read and follow all safety instructions before use.
- Wear protective gloves.
- If exposed or concerned, get medical advice.
- Store locked up.
- Dispose of contents/container in a responsible manner to avoid accidental exposure of uninformed persons and animals.
- Avoid contamination of any food, food utensils and drinking water.
- Set bait where it will be inaccessible to children and animals
- Look for and burn all dead rats and mice

ADDITIONAL MITIGATION MEASURES:

- Ensure the child-resistant re-closable lid is tightly closed at all times.
- Immediately after use, store the container with unused bait in its original container, locked up.
- This remedy may **ONLY** be used in tamper-resistant bait boxes.
- In case of poisoning or suspected poisoning, especially involving children, get immediate medical attention and make this label available.
- Always make sure that baits are adequately protected from children and non-target animals.

RELEVANT SUBSTANCES:

Difenacoum 0.01%

Bittrex < 1%

SYMPTOMS OF POISONING:

Bleeding from the nose and gums; haematura (bleeding under the skin); haemorrhage into and around the elbow and knee joints; blood in faeces and urine. Also symptoms of anaemia, fatigue and dyspnoea.

FIRST AID TREATMENT:

SKIN CONTACT: Wash skin with plenty of soap and cold water.

EYE CONTACT: Rinse eyes with plenty of clean water for at least 15 minutes.

INHALATION: Unlikely to occur with this formulation.

INGESTION: Induce vomiting by tickling the back of the throat. Get medical attention immediately.

Antidote: Vitamin K1 (phytonadione); dosage for adults and children over 12 years – 15-25mg; children under 12 years – 5-10mg. Do not use vitamins K3 or K4 as a substitute for K1.

NOTE TO PHYSICIAN:

Administer vitamin K 1 intravenously depending on prothrombin time on first day. In severe case a small transfusion of fresh blood can be considered.

DIRECTIONS OF USE: Use only as directed.

IMPORTANT:

TORNADO kills after rodents have had one or more feeds. Death will occur 4 to 12 days after a lethal dose has been consumed.

NOTES ON BAITING:

Timber Industry- Newly established plantations.

Place a 20g **TORNADO** wax block at the base of each tree in heavily infested areas. For milder infestations, place at the base of selected trees (16 to 25 blocks/100 trees). Also place near where rodents have nests or burrows. Cover with grass, branches etc. or place in a piece of drain pipe to keep out of view of non- target species.

Industrial Use:

Place 20g **TORNADO** wax blocks at a suitable number of locations near where rodents have nests or burrows or where they have been feeding or cause damage. Inspect bait points regularly and top-up until rodent activity ceases.

Additional Notes:

1. Ensure that domestic and farm animals do not have access to bait by placing it in a piece of drain pipe or under a tile.
2. Replace any baits which have become spoilt or contaminated and dispose of in a responsible manner.
3. Where rodent activity has ceased, remove uneaten bait and destroy in a responsible manner. It is good practice, however to maintain a few permanent feeding points to prevent re-infestation.