

Read the label before opening the container.

For full particulars, see enclosed leaflet.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

BOOSTER AS™

South Africa Reg. No: L6182 Act No. 36 of 1947

ADJUVANT

BOOSTER AS™ is an ammonium sulphate based adjuvant for use with glyphosate based herbicides to improve uptake and aid compatibility. BOOSTER AS™ is classified as non-hazardous according to the Globally Harmonized System (GHS)

ACTIVE INGREDIENT:

Ammonium Sulphate.....500g/ℓ

Product Information: 072 678 8226
In case of poisoning: 082 446 8946

Not classified as hazardous under GHS



Registration holder: RT Chemicals CC

Co. Reg. No: CK87/21925/23

44 Kerk Street, Lichtenburg NW, South-Africa

Tel: +27 12 006 0063

Web: www.envirobiochem.co.za

UN No. Not Regulated

Expiry Date:

Batch No:

Date of Manufacture:



WARNINGS:

- Handle with care.
- Observe all warnings that are applicable to products with which BOOSTER AS™ will be used.
- Store away from food and feedstuffs.
- Keep out of reach of children, uninformed persons and animals.

Although this remedy has been extensively tested under a wide variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instruction or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

- Observe all precautions that are applicable to products with which BOOSTER AS™ will be used.
- Prevent drift of spray mist onto other crops, grazing, rivers, dams or areas not under treatment.
- Clean application equipment after use and dispose of wash water where it will not contaminate crops, grazing, rivers or dams.
- Triple rinse empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the contents of the spray tank before destroying the container in the prescribed manner.
- Empty containers should be taken for local recycling or waste disposal.
- Do not re-use empty containers.
- Prevent contamination of food, feedstuffs, drinking water and eating utensils.

RELEVANT SUBSTANCES:

None known.

FIRST AID TREATMENT:

INHALATION: Remove person to fresh air and keep comfortable for breathing.

INGESTION: Rinse mouth with water. Seek medical attention.

EYE CONTACT: Rinse cautiously with water for several minutes.

DIRECTIONS FOR USE: Use only as directed.

Dosage:

Use 2ℓ BOOSTER AS™ per 100ℓ spray mixture.

Mixing Instructions:

- (a) Half fill the spray tank with clean water, start agitation and add required amount of BOOSTER AS™.
- (b) Add the correct amount of fl owable herbicide, glyphosate and remaining water with continual agitation.
- (c) Should a wetter be required, add this last.
- (d) Maintain adequate agitation during application and do not leave the tank mixture to stand for any length of time.

NOTE: The use of BOOSTER AS™ will increase glyphosate efficacy by improving its uptake and aid compatibility with certain other herbicides.