

Read the label before opening the container.

For full particulars, see enclosed leaflet.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

FOLLIE™ 500

South Africa Reg. No: L11962 Act No. 36 of 1947

FRAC FUNGICIDE GROUP CODE: 4M

A suspension concentrate organic-based fungicide for the preventive control of anthracnose, downy mildew, dead-arm disease and Botrytis rot in wine grapes, dead-arm disease, anthracnose and downy mildew pre-blossom in table grapes as well as late and early blight in potatoes.

ACTIVE INGREDIENT:

Folpet (phtalimide).....500g/ℓ

Product Information: 086 155 5777

In case of poisoning: 082 446 8946

HAZARD STATEMENTS

- May cause an allergic skin reaction
- Causes serious eye irritation
- Harmful is inhaled
- Suspected of causing cancer
- Very toxic to aquatic life

PRECAUTIONARY STATEMENTS

- Store in a well-ventilated place. Keep container tightly closed.
- Use only outdoors or in a well-ventilated area.



DANGER

enviro™
— bio-chem

Registration holder: Enviro Crop Protection (Pty) Ltd

Co. Reg. No: CK 2007/027855/07

44 Kerk Street, Lichtenburg NW, South-Africa

Tel: +27 12 006 0063 | Fax: +27 86 541 7948

Web: www.envirobiochem.co.za

Expiry Date:

Batch No:

Date of Manufacture:

UN No. 3082

WARNINGS:

Withholding periods (minimum number of days between last application and harvest):

Table grapes	No later than pre-blossom stage.
Wine grapes	14 days

NOTE

ABOVE-MENTIONED WITHHOLDING PERIODS REFER TO COMPLIANCE WITH LOCAL MAXIMUM RESIDUE LIMITS (MRL'S). HOWEVER, IT IS IMPORTANT TO NOTE THAT IMPORT TOLERANCES OF OTHER COUNTRIES MIGHT POSSIBLY BE EXCEEDED. IF THE TREATED CROP WILL BE EXPORTED, FIRST CONSULT THE RELEVANT IMPORTER OR EXPORTING BODY REGARDING THE USE OF THIS PRODUCT, MRL'S AND RECOMMENDED WITHHOLDING PERIODS.

- May cause an allergic skin reaction.
- Causes serious eye irritation.
- Harmful is inhaled.
- Suspected of causing cancer.
- Very toxic to aquatic life.
- Keep out of reach of children, uninformed persons, and animals.
- Store in a cool place away from food and feeds.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease to the remedy concerned as well as by the method, time and accuracy of application. The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in event of any uncertainty.

PRECAUTIONS:

- Avoid breathing dust or mist.
- Wash hands, forearms, and face thoroughly after handling. Do not touch eyes.
- Use only outdoors or in a well-ventilated area.
- Contaminated work clothing should not be allowed out of the workplace.
- Avoid release to the environment.
- Wear impervious rubber gloves and boots, protective clothing, and chemical safety goggles.
- Get emergency medical help immediately.
- If exposed or concerned, get medical advice.
- If skin irritation or rash occurs: Get medical advice.
- Take off contaminated clothing and wash it before reuse.
- Store in a well-ventilated place. Keep container tightly closed.
- Store locked up.
- Triple rinse empty container in the following manner: Invert the empty container over the spray or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the content of the spray tank.
- Empty containers should be taken for local recycling or waste disposal.
- Do not re-use empty containers.
- Collect spillage.

- Dispose of contents/container in accordance with local/regional/national regulations.

RELEVANT SUBSTANCES:

Folpet 40%
Antifreeze agent (MEG) <5%

FIRST AID TREATMENT:

Remove patient from source of poisoning and keep him quiet and reassured.

IF SWALLOWED: Immediately call a POISON CENTER/doctor/physician.

Do NOT induce vomiting.

IF ON SKIN: Remove/Take off immediately all contaminated clothing. Rinse skin with water/shower.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do – continue rinsing.

IF INHALED: Remove person to fresh air and keep comfortable for breathing.

DIRECTIONS FOR USE: Use only as directed.

Compatibility:

Follie™ 500 can be applied in a tank-mix with most of the commonly used insecticides and fungicides with the exception of Lime, Lime Sulphur and Bordeaux mixture. It should also not be applied with any other *Botrytis* remedy in wine and table grapes.

Application:

Apply as a preventive application and ensure thorough coverage of both upper and lower leaf surface.

APPLICATION RATES:

CROP/DISEASE	DOSAGE	REMARKS
POTATOES Late blight (<i>Phytophthora infestans</i>) Early blight (<i>Alternaria solani</i>)	2ℓ/ha	Preventive treatment: Apply from emergence at 7- to 10-day intervals in not less than 500ℓ water/ha in the pre-blossom stage and 1000ℓ/ha in the post-blossom stage. For the control of early and late blight, particularly after flowering, it is important to obtain thorough coverage. A suitable wetter can be used in the spray mixture.
TABLE GRAPES Deadarm disease, anthracnose, downy mildew	200mℓ/100ℓ water	Recommended only until pre-blossom stage. Apply first spray at bud swell stage. Repeat at 7-to 10-day intervals depending on weather conditions.

CROP/DISEASE	DOSAGE	REMARKS
WINE GRAPES Anthracnose, deadarm disease and <i>Botrytis</i> rot Downy mildew	 200mℓ/100ℓ water 200mℓ/100ℓ water	 Start application at bud swell. Repeat weekly when weather conditions promote disease development. Pre-blossom: First application when shoots are 10 cm long. Repeat at 7- to 10-day intervals depending on weather conditions. Blossom to 80 % petal drop: Apply only when blossoming period lasts for more than 14 days. Post blossom: Apply twice or more at 7- to 14-day intervals depending on weather conditions.